



Cite this article: Bongaerts P, Carmichael M, Hay KB, Tonk L, Frade PR, Hoegh-Guldberg O. 2015 Prevalent endosymbiont zonation shapes the depth distributions of scleractinian coral species. *R. Soc. open sci.* **2**: 140297. <http://dx.doi.org/10.1098/rsos.140297>

Received: 10 September 2014

Accepted: 13 January 2015

Subject Category:

Genetics

Subject Areas:

ecology/evolution

Keywords:

coral reef, depth distribution, symbiont zonation, mesophotic, *Symbiodinium*

Author for correspondence:

Pim Bongaerts

e-mail: pim@uq.edu.au

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsos.140297> or via <http://rsos.royalsocietypublishing.org>.

Prevalent endosymbiont zonation shapes the depth distributions of scleractinian coral species

Pim Bongaerts^{1,2,3}, Margaux Carmichael¹, Kyra B. Hay¹, Linda Tonk^{1,2}, Pedro R. Frade^{3,4} and Ove Hoegh-Guldberg^{1,2}

¹Global Change Institute, and ²ARC Centre of Excellence for Coral Reef Studies, The University of Queensland, St Lucia, Queensland 4072, Australia

³CARMABI, Piscaderabaai z/n, PO Box 2090, Willemstad, Curaçao

⁴Department of Limnology and Bio-Oceanography, Division of Marine Biology, University of Vienna, Althanstrasse 14, Vienna 1090, Austria

1. Summary

Bathymetric distributions of photosynthetic marine invertebrate species are relatively well studied, however the importance of symbiont zonation (i.e. hosting of distinct algal endosymbiont communities over depth) in determining these depth distributions still remains unclear. Here, we assess the prevalence of symbiont zonation in tropical scleractinian corals by genotyping the *Symbiodinium* of the 25 most common species over a large depth range (down to 60 m) on a Caribbean reef. Symbiont depth zonation was found to be common on a reef-wide scale (11 out of 25 coral species), and a dominant feature in species with the widest depth distributions. With regards to reproductive strategy, symbiont zonation was more common in broadcasting species, which also exhibited a higher level of polymorphism in the symbiont zonation (i.e. number of different *Symbiodinium* profiles involved). Species with symbiont zonation exhibited significantly broader depth distributions than those without, highlighting the role of symbiont zonation in shaping the vertical distributions of the coral host. Overall, the results demonstrate that coral reefs can consist of highly structured communities over depth when considering both the coral host and their obligate photosymbionts, which probably has strong implications for the extent of connectivity between shallow and mesophotic habitats.

2. Introduction

Tropical scleractinian corals have been documented across a wide bathymetric range (0 to more than 100 m) within the photic

zone [1], with coral species exhibiting distinct vertical distributions [2–4]. The strong environmental gradients encountered across this bathymetric range are important drivers in this zonation, particularly factors such as wave action, heterotrophic resource availability and temperature as they can differ greatly at opposite ends of the depth spectrum [4–7]. However, as scleractinian corals depend to a great extent on photosynthesis for their energy requirements due to their obligate symbiosis with dinoflagellates from the genus *Symbiodinium* [8], the exponential decrease in available irradiance probably plays one of the most important roles in determining the vertical distributions of coral species [4]. Coral species exhibit adaptations along this light gradient to optimize light capture by their endosymbionts, ranging from changes in gross colony morphology [9], skeletal morphology [10], to host pigment composition [11]. In addition, coral populations of the same species at opposite ends of their depth range have been found in association with different *Symbiodinium* types (e.g. [6,12–14]) that can be physiologically distinct and adapted to different light conditions [15,16]. Such ‘symbiont zonation’ (i.e. a bathymetric shift in associated *Symbiodinium*) has been observed for a range of different coral species (e.g. [6,12–14]), and probably facilitates distribution of these species across a broad environmental gradient [16]. However, the prevalence of ‘symbiont zonation’ on a reef-wide scale remains unknown, and it is unclear to what extent it underlies the vertical distributions of coral species.

Most assessments of *Symbiodinium* diversity over depth have been focused either on several species (reviewed in [17]), or community-wide but with low numbers of replicate samples to allow for extensive taxon sampling [18–21]. In addition, most *Symbiodinium* studies have been carried out over narrow depth spans, often sampling target species over a subsection of their entire depth distribution (but see [6,7,22–24]), limiting the power of detecting symbiont zonation. While depth zonation in associated *Symbiodinium* has been observed for coral species with both narrow [12,14] and broad [6,7] depth distributions, it remains unclear whether there is a link between the vertical distribution of the coral host and the occurrence of symbiont zonation (e.g. whether symbiont zonation is more common in corals with broader depth distributions). Similarly, endosymbiont zonation has been found in both brooding and broadcasting coral species [17], but it is uncertain whether *Symbiodinium* zonation is more common in either reproductive mode. Offspring from broadcasting corals generally acquire *Symbiodinium* from the water column (horizontal symbiont transmission), whereas most brooding corals are thought to acquire *Symbiodinium* directly from the maternal colony (horizontal symbiont transmission) [25,26]. This difference in endosymbiont acquisition mode could potentially lead to distinct patterns of *Symbiodinium* diversity over depth [27]. However, assessing links between host traits (i.e. depth distribution and reproductive mode) and patterns of *Symbiodinium* over depth requires extensive genotyping across both a wide number of species and a broad bathymetric range.

The diversity of *Symbiodinium* at mesophotic depths remains poorly explored, and as such, the extent of specialization of mesophotic endosymbiont communities remains unclear [27]. Initial studies in the Caribbean identified putative ‘deep-specialist’ *Symbiodinium* types in the genera *Madracis* [6], *Montastraea* [7] and *Agaricia* [24], pointing towards a certain level of endosymbiont specialization at mesophotic depths. By contrast, preliminary studies on the Great Barrier Reef and in Hawai’i have indicated mesophotic corals associate predominantly with *Symbiodinium* types commonly found in shallow water [27,28]. Nonetheless, most studies of *Symbiodinium* diversity have been carried out over relatively shallow depth ranges, and it may well be that specialization only occurs deeper into the mesophotic zone. In terms of coral species distributions [17,29], the upper mesophotic appears to represent a transition zone between the shallow and lower mesophotic [29,30], and a similar pattern may be present for the associated endosymbiont community. However, the extent of specialization of mesophotic *Symbiodinium* communities and the overlap of endosymbiont communities between shallow and mesophotic depths remains largely untested.

The wide bathymetric distribution of certain coral species, and the relatively shallow depth to which certain stressors and disturbances are limited, has led to the hypothesis that deep reefs may act as refugia and provide a source of reproduction for their shallow-water counterparts (reviewed in [17]). However, recent molecular ecological studies have demonstrated that corals with broad depth distributions do not necessarily form a single metapopulation [23,30–34], and the ability of a species to act as a source of reproduction should therefore not be based on species distribution alone (i.e. co-occurrence in shallow and deep habitats). The relative difficulties associated with the development of informative genetic markers [35] and, more recently, genotyping-by-sequencing approaches for cnidarians hosting endosymbionts still hamper assessment of host population structure on an ecosystem-wide scale (i.e. across a wide range of species), and genotyping efforts have therefore been restricted to case studies of particular species. However, several studies have found that host populations that harbour distinct *Symbiodinium* types over depth (e.g. *Seriatopora hystrix*, *Madracis pharensis* and *Montastraea cavernosa*) are




Figure 1. (a) Bathymetric profile at sampling location (Buoy 0/1, Curaçao), (b) number of coral host species and associated *Symbiodinium* profiles observed across the sampled depth range (2–60 m) and (c) prevalence of clades and clade combinations over depth in the *Symbiodinium* profiles of sampled specimens (with numbers of sampled colonies indicated).

genetically divergent [30–33], illustrating that the occurrence of symbiont zonation can be an indication or ‘proxy’ of underlying host genetic structuring. As such, assessment of symbiont zonation across a broad range of species (which can be assessed using a single *Symbiodinium* marker) may be an important first step towards a better understanding of the potential extent of host population structuring over depth and vertical connectivity of coral communities on a reef-wide scale.

Although we have a reasonable understanding of the vertical distribution of invertebrate species on coral reefs, the role of symbiont zonation underlying these distributions remains poorly understood, particularly on an ecosystem-wide scale. Here, we assess *Symbiodinium* diversity over a large depth range (2–60 m) and across 25 coral species to determine: (i) the overall prevalence of symbiont depth zonation and (ii) whether the occurrence of symbiont zonation is linked to reproductive mode and/or depth distribution of the host. *Symbiodinium* identity was established for 16 broadcasting and nine brooding coral species across depth intervals (2, 5, 10, 25, 40, 50 and 60 m) for a total of 1636 coral colonies from the Buoy 0/1 study site in Curaçao, using ITS2/DGGE fingerprinting. Representing one of the largest genotyping efforts carried out on a single reef, this study demonstrates that coral reefs can consist of highly structured communities when taking into account both the coral host and its associated *Symbiodinium*.

3. Material and methods

3.1. Sample collection

Coral samples were collected at the Buoy 0/1 site situated 500 m west of the CARMABI Institute, Curaçao, Southern Caribbean (12°07′31 N, 68°58′27 W) (figure 1). The study site is characterized by a 50–100 m wide shallow terrace that gently slopes down to 8–12 m, followed by a fore reef slope that shows an abrupt descent down to a 50–60 m deep terrace (figure 1a). Colonies were sampled from 25 coral species at this study site, with the majority of samples being collected during the following sampling periods: July–August 2008, November 2009 and the ‘Catlin Seaview Survey’ expedition in March–April 2013. We also included samples from two previously published datasets of the genus *Madracis* ($n = 307$; collected between 2004 and 2006) [6,16,32] and *Agaricia* ($n = 332$; collected between 2004 and 2009) [24] that were obtained at the exact same study site. Care was taken to avoid resampling of colonies in different sampling years. Small coral fragments (approx. 3 cm²) were collected by hammer and chisel at the following depth intervals: 2 m (± 1 m), 5 m (± 1 m), 10 m (± 1 m), 25 m (± 2 m), 40 m (± 2 m), 50 (± 2 m) and 60 m (± 2 m) for each species that could be found at that depth. Species identifications were done *in situ*, and later reconfirmed in the laboratory before processing. Coral tissue samples were stored in ethanol or a salt-saturated 20% DMSO/0.5 M EDTA solution at -20°C . Total DNA was extracted from the tissue using a Qiagen Plant Mini Kit, MoBio Ultra Clean Soil DNA Kit (following the manufacturer’s instructions), or a slightly modified method used for black tiger shrimp [23,36].

3.2. Genetic characterization of associated algal endosymbionts (*Symbiodinium*)

The internal transcribed spacer 2 (ITS2) region of the rDNA for *Symbiodinium* was amplified for all samples ($n = 1636$) using *Symbiodinium*-specific primers ([18]) as described in Bongaerts *et al.* [31]. To identify the dominant *Symbiodinium* types in each sample, the amplified ITS2 fragments were separated using denaturing gradient gel electrophoresis (DGGE) on a CBScientific System following conditions outlined in Sampayo *et al.* [14]. Running conditions varied for the *Madracis* samples (i.e. 30–70% gradient on Bio-Rad DCode system) included from Frade *et al.* [6], however representative newly acquired *Madracis* samples were run on both systems to allow for cross-comparison between DGGE profiles. Representative, dominant bands of each characteristic profile were excised (usually from several replicate profiles), eluted overnight in dH₂O, re-amplified and purified (using ExoSAP-IT) prior to sequencing. The re-amplified PCR products were sequenced in both the forward and reverse directions (ABI BigDye Terminator chemistry, Australian Genome Research Facility). Chromatograms were analysed using CodonCode aligner with sequences being aligned with MUSCLE and blasted on GenBank (<http://www.ncbi.nlm.nih.gov/BLAST>). A haplotype network was created using TCS 1.21 [37], treating gaps as a fifth character state. Network analyses were performed separately for *Symbiodinium* clade B and C because of their high level of divergence. Coral–*Symbiodinium* associations were compared to previously described associations using the GeoSymbio [38] and SD2-GED [39] databases.

The co-dominant sequences in a given profile may represent either intragenomic variants and/or a mix of distinct *Symbiodinium* types, however as this inference can sometimes be subjective we named profiles in this study by their co-dominant ITS2 ‘sequences’ (e.g. C3/C3b/C3nN1) rather than symbiont ‘types’ (which can consist of multiple sequences). Known *Symbiodinium* sequences are indicated by their ITS2 sequence name *sensu* LaJeunesse (e.g. C3d), novel sequences are identified by a nomenclature that specifies the known sequence to which they are most related, followed by a capital N (indicating novel sequence) and an arbitrary number (e.g. C3b.N1) and incomplete sequences (i.e. shorter length) are identified by a nomenclature that specifies the known sequence to which they are most related (based on their partial sequence) followed by a capital U (indicating full sequence remains ‘unknown’) and an arbitrary number (e.g. C3.U1).

3.3. Mitochondrial DNA sequence analyses of coral colonies

The coral host mitochondrial *nad5* non-coding region was amplified using primers F18 (5'-GTCCTTACGTCTTTACACCGAC-3') and R17 (5'-AAAGACCACTCTAAAGCCCCGT-3') for randomly picked samples from the shallowest and deepest collection depth of each of the coral species in this study ($n = 90$) to check for potential cryptic diversity across the sampled bathymetric range (as done for *Madracis* spp. previously by Frade *et al.* [32]). PCR amplifications were performed using conditions described for the *atp6* gene in Bongaerts *et al.* [24], and products were purified, sequenced, aligned and analysed as specified for the algal endosymbionts (*Symbiodinium*). Phylogenetic analyses of sequences were performed using maximum likelihood in MEGA 6 [40] under the delayed transition setting and calculation of bootstrap support values based on 1000 replicates. The best-fit model of molecular evolution was selected by hierarchical Akaike information criterion (AIC) using MEGA 6 [40] with a HKY + G model best describing the *nad5* data under a log likelihood optimality criterion.

3.4. Statistical analyses

For host species associating with more than one *Symbiodinium* profile, an (nested) analysis of similarity (ANOSIM) was carried out using a Sorensen resemblance measure in the software package PRIMER v. 6 to test for differences between sampling years (two-way ANOSIM with depth nested within year) and depths (one-way ANOSIM comparing individual depth groups). Data from depths where less than 10 coral colonies were sampled were not included in the ANOSIM. Fisher's exact tests (two-tailed P) were used to investigate the relation between reproductive mode and the occurrence of symbiont zonation, and the relation between reproductive mode and the level of polymorphism observed in symbiont zonation (two to three profiles or more than three profiles) using numbers of species as values. A Mann–Whitney U -test was used to assess differences in *Symbiodinium* diversity between broadcasting and brooding species. The effect of symbiont zonation and reproductive mode on the depth distribution of the host (determined as the difference between the shallowest and deepest sampling depth) was assessed using two-way ANOVA, with data $\log(x + 10)$ transformed to meet the assumptions of normality and homoscedasticity.

4. Results

4.1. *Symbiodinium* diversity and host–symbiont associations

Symbiodinium community profiles were assessed for 1636 coral colonies originating from a single reef, of which 375 originated from mesophotic depths (40–60 m). In total, 93 *Symbiodinium* DGGE profiles were distinguished across the 25 studied coral species (figure 1b), based on a total of 120 diagnostic bands. A large amount of profiles only consisted of a single dominant band (41%), however others consisted of either multiple bands from the same *Symbiodinium* clade (24%) or multiple bands belonging to different *Symbiodinium* clades (35%) (figure 2). These diagnostic bands represented 22 known *Symbiodinium* sequences and 39 novel *Symbiodinium* sequences (electronic supplementary material, figure S1) (GenBank accession nos. KP178779–KP178809, KF551185–KF551192), with 22 bands yielding incomplete sequences (after multiple sequencing attempts). Most obtained *Symbiodinium* sequences belonged to either clade C (61%) or clade B (32%), with some belonging to the A, D and H clades which were found exclusively in shallow habitats (less than 3% each) (figure 1c).

Several of the observed *Symbiodinium* ITS2 sequences in this study have been observed previously for the same species in other Caribbean locations, such as B7 and B13 in *Madracis mirabilis*, B7 in *Madracis pharensis*, D1a in *Colpophyllia natans*, C3 in *Diploria strigosa* and C12 in *Orbicella faveolata* [38,39]. However, for many species we observed sequences (in both shallow and deep water) that have not been previously identified in association with those species, ranging from closely related variants (e.g. C16.N16 in *Stephanocoenia intersepta*) to distinct clades (e.g. D1a in *Mycetophyllia ferox*) (figure 2). Coral species with a brooding reproductive mode harboured significantly lower numbers of *Symbiodinium* profiles than species with a broadcasting reproductive mode (Mann–Whitney *U*-test, $U = 13$, $Z = -3.4$, $p < 0.001$). An exception was *Porites astreoides* that hosted a large number of *Symbiodinium* profiles, containing many novel ITS2 sequences. Broadcast-spawning species harboured higher symbiont diversity and many ITS2 sequence variants were identified that were quite divergent from types previously described for those species (e.g. C7 and C16 in *Siderastrea siderea*) (figure 2). Additionally, broadcasting species frequently harboured symbionts from at least two different clades, whereas this was uncommon in brooding species.

4.2. Coral host identity

We sequenced the *nad5* region for a subset of individuals ($n = 90$) as an initial verification to assess whether shallow and deep populations of collected coral species indeed represent the same ‘species’ (i.e. are not genetically divergent for a conserved DNA region). Lengths of the sequenced *nad5* region varied between 621 and 911 bp depending on whether the coral species belonged to, respectively, the robust or complex phylogenetic group, and sequences were aligned separately based on these groupings (GenBank accession nos. KP178822–KP178861, KP178862–KP178900). Most species (15 out of 19) consisted of a single haplotype per species (with no difference between shallow and deep individuals), and the three species consisting of multiple haplotypes (*C. natans*, *S. intersepta* and *Agaricia lamarcki*) showed no clear partitioning between shallow and deep individuals (i.e. haplotypes were shared) (electronic supplementary material, figure S2). The *nad5* marker was too conserved to distinguish between Faviidae (e.g. *Colpophyllia* and *Diploria*) and members of the *Orbicella* species complex (*O. annularis*, *O. faveolata* and *O. franksi*). Divergence between *Madracis* species was assessed in a previous study [32] that reported genetic divergence between shallow and deep populations of *M. pharensis* at this sampling location (based on the *nad5* marker and two nuclear markers).

4.3. Prevalence of symbiont zonation

Depth distributions of individual coral species varied, with absence of *Symbiodinium* data for a species at a particular sampling depth meaning that the species was not observed at that depth during the more than 75 collection dives (conducted to collect all the specimens) on the study site (figure 2). Eleven species with varying depth distributions (Δ depth = 10–55 m) exhibited significant differences in *Symbiodinium* community between sampling depths (i.e. symbiont zonation) (figure 2; electronic supplementary material, table S1). In four cases this consisted of a shift between only two and three *Symbiodinium* profiles (*A. lamarcki*, *M. mirabilis*, *M. pharensis* and *M. formosa*), and in the other cases consisted of a more complex shift (i.e. involving more than three *Symbiodinium* profiles). By contrast, eight species associated with a single discernable *Symbiodinium* profile across their entire depth distribution (Δ depth = 0–45 m) and five species associated with several different *Symbiodinium* profiles but with




Figure 3. Mean depth distribution ranges for broadcasting and brooding species with and without symbiont zonation. Error bars indicate s.e.m.

no significant differences between depth groups (figure 2). One species (*Madracis carmabi*) was only found at one sampling depth, but hosted a singled *Symbiodinium* profile. No significant differences were found between sample groups collected during different fieldwork periods (two-way ANOSIM with depth nested within year) (electronic supplementary material, table S1).

Four out of the five species with the largest depth distributions (Δ depth = 40–55 m; *M. pharensis*, *S. siderea*, *S. intersepta* and *M. cavernosa*) exhibited significant symbiont zonation (with *Agaricia agaricites* being the exception), whereas only one of the nine species (*O. franksi*) with the smallest depth distributions (Δ depth \leq 15 m) exhibited symbiont zonation (figure 2). With regards to reproductive strategy, only five out of 16 brooding coral species exhibited symbiont zonation, while for broadcast-spawning species more than half did (six out of nine), however there was no significant association (Fisher's exact test, $p > 0.05$) between reproductive strategy and the occurrence of *Symbiodinium* zonation. Symbiont zonation in all broadcasting coral species was highly polymorphic (i.e. involved more than three *Symbiodinium* profiles), whereas in all brooding species the zonation was 'simple' (i.e. involved two to three *Symbiodinium* profiles), with the exception of *P. astreoides*, and this association was significant (Fisher's exact test, $p < 0.05$). Species that exhibited symbiont zonation had significantly larger depth distributions (two-way ANOVA, $F_{1,21} = 10.0$, $p < 0.005$), whereas there was no significant effect of reproductive mode or interaction between reproductive mode and symbiont zonation (figure 3) (electronic supplementary material, table S2).

4.4. Mesophotic *Symbiodinium* diversity

Compared with shallow depths (5–25 m), the diversity of scleractinian corals and associated *Symbiodinium* diversity (both in terms of absolute numbers and numbers of profiles per species) at mesophotic depths (40–60 m) was substantially lower (figure 1b). Depth-generalist species that occurred down to 40–50 m depth mostly harboured the same *Symbiodinium* profile as in shallower water (10–25 m), however depth-generalist species that extend down to 60 m depth (i.e. *S. intersepta*, *M. cavernosa* and *M. pharensis*) all exhibited distinct *Symbiodinium* profiles at their deeper extremes (figure 2). Nonetheless, the common symbiont profiles associated with *S. intersepta* at 60 m represent *Symbiodinium* types (C1 and C3) that commonly occur on shallow reefs (but not in this species at the study location). Depth-'breakpoints' were assessed by evaluating pairwise differences between depths for each species (electronic supplementary material, table S1), with 'breakpoints' indicating where *Symbiodinium* communities from all depths above and below are significantly different from each other (e.g. for *A. lamarcki* the 'breakpoint' is located between 25 and 40 m, as the *Symbiodinium* community at 10 m = 25 m \neq 40 m = 50 m). The depth at which these 'breakpoints' occurred varied between species, however they were most frequently observed between 10 and 25 m ($n = 6$), with only one breakpoint observed at shallower depths (i.e. between 2 and 5 m). The remainder occurred at the intersection of the mesophotic zone (between 25 and 40 m; $n = 2$) or within the mesophotic zone (between 40 and 50 m or 50 and 60 m; $n = 5$).

5. Discussion

Depth zonation in associated *Symbiodinium* has been demonstrated for individual scleractinian coral species and genera, with populations hosting distinct endosymbionts at different ends of the depth spectrum (e.g. [12,13,16,41,42]). Here, we demonstrate through extensive genotyping that such symbiont zonation is in fact common among scleractinian corals, with nearly half of the studied species showing significant changes in their *Symbiodinium* community over depth (figure 2). In addition, we found that species exhibiting symbiont zonation have significantly wider depth distributions, which is particularly apparent in broadcasting species (figure 3). We conclude that endosymbiont zonation plays an important role in determining the bathymetric distribution of Caribbean corals, with its overall prevalence leading to highly structured coral host–endosymbiont communities over depth.

Overall symbiont diversity was found to be high, corroborating previous studies in other Caribbean regions [18,21], with only a quarter of the obtained sequences representing known types that have been observed in other parts of the Caribbean. It also corroborates the dominance of clade B and C in this region, with the occurrence of other clades (A, D and H) being restricted almost exclusively to shallow water (10 m or less) [18,21] (figure 1c). The lower *Symbiodinium* diversity at mesophotic depths (40–60 m) is in line with previous observations of decreasing diversity over smaller depth ranges in the Caribbean [18,21]. Many new host–*Symbiodinium* associations were observed, with most coral species harbouring profiles containing *Symbiodinium* sequences (ranging from closely related variants to distinct clades) that have not been previously identified in association with those species (figure 2). Observing novel host–*Symbiodinium* associations is common when surveying new geographical regions [43], mesophotic depths [27] or when increasing sample sizes [14], and demonstrates the importance of expanded surveying to broaden our understanding of specificity and diversity in the coral–algal symbiosis. The many novel associations (and ITS2 sequence types) observed here highlight the extensive diversity underlying the geographical and ecological distribution of Caribbean corals.

Symbiont zonation was observed in four out of five coral species with the widest depth distributions, indicating that a change in *Symbiodinium* over depth is a common trait associated with ‘depth-generalist’ coral species. The association with different endosymbionts over depth probably enables these species to thrive under the different environmental conditions encountered along their broad depth distribution ranges [16], through adaptation to particular light [15], temperature [44,45] or potentially other environmental conditions such as light spectrum or nutrient availability [16]. Nonetheless, the fact that *Agaricia agaricites* associated with a single *Symbiodinium* profile along its entire bathymetric distribution (5–50 m) (figure 2) demonstrates that symbiont zonation is not an obligate strategy for depth-generalist species [24]. A similar lack of symbiont zonation has been observed for the Indo-Pacific coral species *Pachyseris speciosa*, which hosted predominantly a single *Symbiodinium* profile despite its broad depth distribution in northwest Australia (10–60 m) [22]. While the observed patterns point towards symbiont zonation being more common in species with large depth distributions, symbiont zonation was also observed over very small depth ranges (e.g. in *Orbicella franski*: Δ depth = 15 m), which is not entirely surprising as certain environmental gradients (e.g. incident irradiance, ultraviolet light and wave action) are most pronounced across shallow depth ranges. Given the inability of ITS2-DGGE to identify *Symbiodinium* at background levels [46] and the fact that only dominant bands are used for profile assignment, assessment of changes in associated *Symbiodinium* diversity over depth remains conservative as it ignores subtle changes in relative *Symbiodinium* abundances. As such, it could be that the frequency of symbiont zonation remains underestimated, however this would not affect the conclusion that *Symbiodinium* zonation is common on a reef-wide scale.

Clear differences were observed between reproductive strategies in the diversity of associated *Symbiodinium*, with broadcasters always associating with multiple *Symbiodinium* profiles (sometimes representing different clades) and with patterns of symbiont zonation always involving at least four (but often more) different *Symbiodinium* profiles. By contrast, brooders often associated with a single *Symbiodinium* profile, and when exhibiting symbiont zonation this usually only involved two to three distinct profiles representing a single clade (with *P. astreoides* as exception). At first, these differences seem to be associated with the different symbiont acquisition modes (respectively, horizontal and vertical) commonly associated with broadcasting and brooding species [25]. Indeed, the broadcasting species studied here reportedly acquire their *Symbiodinium* horizontally [34,47] (figure 2), and the higher *Symbiodinium* diversity and prevalence of symbiont zonation may be facilitated by the ‘open nature’ of the coral–algal symbiosis representative of this acquisition mode [48–50]. The many novel associations that were observed for these species may be a further consequence of this potential ‘flexibility’, and highlights the potential role this acquisition mode has played in the ecological and

geographical range extensions of these coral species. By contrast, many horizontally transmitting species in the Indo-Pacific appear to associate with relatively few *Symbiodinium* types [20,51,52], which in part may be a consequence of the overall lower *Symbiodinium* diversity in this region compared with the Caribbean [19].

Brooding species appear to exhibit a variety of *Symbiodinium* acquisition strategies, including vertical (e.g. *Agaricia humilis*), horizontal (e.g. *Eusmilia fastigiata*) and facultative symbiont transmission (e.g. *Madracis mirabilis*) [47] (figure 2). In addition, a recent study reports that *Symbiodinium* can sometimes be acquired through both horizontal and vertical symbiont acquisition (e.g. *Stylophora pistillata* [50]). Thus, while the transmission mode remains unclear for some of the brooding species, the observed pattern is that of low diversity on a localized scale, in line with the relatively 'closed nature' of a vertical symbiont acquisition mode. In a vertical symbiont acquisition mode, different *Symbiodinium* associations are thought to arise through (co-)evolutionary processes and/or (relatively rare) host-symbiont recombination events [24,51] that can become fixed through ecological selection or geographical isolation. Such processes are probably responsible for the novel associations observed for these brooding species, and more broadly for geographical differences in *Symbiodinium* diversity of brooding species, rather than merely reflecting symbiotic 'flexibility'. Similarly, on the Great Barrier Reef, where initially diverse associations were described for several vertically transmitting pocilloporid species (e.g. [14,52]), it later became apparent that these 'species' in fact comprise multiple lineages/species with specific endosymbiont associations (e.g. [31,53]). Nonetheless, while *Symbiodinium* diversity associated with individual brooding species is often low on a localized scale, the overall diversity can in fact be high due to biogeographic differences, with higher levels of diversity compared with horizontally transmitting species [54]. The brooding species *Porites astreoides* represents an exception with its extremely high local *Symbiodinium* diversity and indicates that *Symbiodinium* acquisition in this brooding species may not be limited to only maternal acquisition [55], and potentially represents a more 'open' symbiosis.

The associated endosymbionts at the lower reaches of this study (50–60 m) appear to represent a specialized community, consisting mostly of *Symbiodinium* that are not observed in shallow water (except, for example, those associated with *A. agaricites*) (figure 2), which is in contrast with preliminary observations from the Indo-Pacific region [27,28]. For some species (e.g. *Stephanocoenia intersepta* and *Montastraea cavernosa*), the symbiont community was distinct even between different depths within the mesophotic zone, which has been observed previously for *M. cavernosa* in the Bahamas [7,33]. Interestingly, the deepest *S. intersepta* populations (60 m) associated at our study site with the generalist types *Symbiodinium* C1 and/or C3 that are commonly found in shallow water [56]. The presence of these *Symbiodinium* at mesophotic depths could be interpreted as a lack of a specialized endosymbiont community, however the absence of these types within the shallow *S. intersepta* populations at our study site points towards (local) specialization of this host-*Symbiodinium* association to mesophotic conditions (60 m). As such, ecological implications of 'depth-specialization' should therefore be assessed by looking at the combination of symbiotic partners and on a localized scale (i.e. by comparing shallow versus deep assemblages at a single location). Alternatively, it may be that these *Symbiodinium* are physiologically distinct through recent adaptation to different environmental conditions, as adaptive variation has been demonstrated in *Symbiodinium* C1 previously [57]. At upper mesophotic depths (40 m) of the study site, most of the dominant *Symbiodinium* profiles can be found also at shallower depths for the same species. This indicates that there is substantial overlap in *Symbiodinium* associations between the upper reaches of the mesophotic zone and intermediate/shallow depths for depth-generalist coral species, and reinforces the idea of the upper mesophotic as a transition zone between the shallow and lower mesophotic reef [29,30].

Symbiont zonation has been found to be associated with genetic structuring of the coral host in several brooding and a broadcasting species [7,31–34], and as such can be indicative of underlying host genetic differentiation. The patterns of *Symbiodinium* observed here on a reef-wide scale might therefore be reflective of highly structured coral host populations over depth and genetically distinct populations at lower mesophotic depths. On the other hand, there is considerable overlap in host-*Symbiodinium* associations observed between shallow (10–25 m) and upper mesophotic depths (40 m), highlighting the potential connectedness of these habitats. Nonetheless, host genetic differentiation across habitats has also been observed between populations hosting the same *Symbiodinium* type [31,58], and conversely a lack of host differentiation has been observed for species exhibiting symbiont zonation [59]. Our initial screening of host genetic differentiation was aimed at identifying potential cryptic species diversity and did not reveal any gross genetic differences, other than the depth-divergence of *Madracis pharensis* that was reported in Frade *et al.* [32]. However, the targeted mitochondrial region (*nad5*) was too conserved to assess population-level genetic differentiation (and within the robust clade to assess even intrageneric

differences), and the presence of host genetic differentiation underlying the *Symbiodinium* zonation remains to be assessed. The prevalence of *Symbiodinium* zonation stresses the importance of such future host-focused studies, in particular in assessing reef-wide patterns of vertical connectivity.

In conclusion, this study demonstrates the prevalence of endosymbiont zonation underlying the vertical distributions of Caribbean scleractinian corals and indicates that *Symbiodinium* zonation appears to play an important role in facilitating the broad depth ranges of depth-generalist species. Future studies should assess whether this pattern is consistent with other geographical regions and whether endosymbiont zonation is also pervasive in other non-scleractinian photosynthetic marine invertebrates. Overall, we clearly need a better understanding of the ecological and evolutionary interactions between coral hosts and their endosymbionts, through high-resolution genetic studies of both symbiotic partners, to further elucidate reef-wide patterns of host-*Symbiodinium* structuring and the interconnectedness of habitats on coral reefs.

Data accessibility. Sequence data have been deposited in GenBank under the following accession nos. KP178779–KP178809, KP178822–KP178900.

Acknowledgements. We would like to thank Mark Vermeij, Norbert Englebert, Carlos Winterdaal, Judith Bakker and Joshua Boldt for logistical support.

Funding statement. The study was conducted as part of the ‘Catlin Seaview Survey’ funded by Catlin Group Limited, with support from the Australian Research Council (ARC) Centre of Excellence for Coral Reef Studies and the Global Change Institute.

References

- Kahng SE, Garcia-Sais JR, Spalding HL, Brokovich E, Wagner D, Weil E, Hinderstein L, Toonen RJ. 2010 Community ecology of mesophotic coral reef ecosystems. *Coral Reefs* **29**, 255–275. (doi:10.1007/s00338-010-0593-6)
- Goreau TF, Wells JW. 1967 The shallow-water Scleractinia of Jamaica: revised list of species and their vertical distribution range. *Bull. Mar. Sci.* **17**, 442–453.
- Sheppard C. 1982 Coral populations on reef slopes and their major controls. *Mar. Ecol. Prog. Ser.* **7**, 83–115. (doi:10.3354/meps007083)
- Veron JEN, Stafford-Smith M. 2000 *Coral reefs of the world*. Townsville, Australia: Australian Institute of Marine Science.
- Dollar SJ. 1982 Wave stress and coral community structure in Hawaii. *Coral Reefs* **1**, 71–81. (doi:10.1007/BF00301688)
- Frade PR, De Jongh F, Vermeulen F, van Bleijswijk J, Bak RPM. 2008 Variation in symbiont distribution between closely related coral species over large depth ranges. *Mol. Ecol.* **17**, 691–703. (doi:10.1111/j.1365-294X.2007.03612.x)
- Lesser MP, Slattery M, Stat M, Ojimi M, Gates RD, Grotto A. 2010 Photoacclimatization by the coral *Montastraea cavernosa* in the mesophotic zone: light, food, and genetics. *Ecology* **91**, 990–1003. (doi:10.1890/09-0313.1)
- Trench RK. 1993 Microalgal-invertebrate symbioses—a review. *Endocytobios. Cell Res.* **9**, 135–175.
- Titlyanov EA. 1987 Structure and morphological differences of colonies of reef-building branched corals from habitats with different light conditions. *Mar. Biol.* **1**, 32–36.
- Enriquez S, Méndez ER, Iglesias-Prieto R. 2005 Multiple scattering on coral skeletons enhances light absorption by symbiotic algae. *Limnol. Oceanogr.* **50**, 1025–1032. (doi:10.4319/lo.2005.50.4.1025)
- Salih A, Larkum A, Cox G, Kühl M, Hoegh-Guldberg O. 2000 Fluorescent pigments in corals are photoprotective. *Nature* **408**, 850–853. (doi:10.1038/35048564)
- Rowan R, Knowlton N. 1995 Intraspecific diversity and ecological zonation in coral-algal symbiosis. *Proc. Natl Acad. Sci. USA* **92**, 2850–2853. (doi:10.1073/pnas.92.7.2850)
- Warner ME, LaJeunesse TC, Robison JD, Thur RM. 2006 The ecological distribution and comparative photobiology of symbiotic dinoflagellates from reef corals in Belize: potential implications for coral bleaching. *Limnol. Oceanogr.* **51**, 1887–1897. (doi:10.4319/lo.2006.51.4.1887)
- Sampayo EM, Franceschinis L, Hoegh-Guldberg O, Dove S. 2007 Niche partitioning of closely related symbiotic dinoflagellates. *Mol. Ecol.* **16**, 3721–3733. (doi:10.1111/j.1365-294X.2007.03403.x)
- Iglesias-Prieto R, Beltrán VH, LaJeunesse TC, Reyes-Bonilla H, Thomé PE. 2004 Different algal symbionts explain the vertical distribution of dominant reef corals in the eastern Pacific. *Proc. R. Soc. Lond. B* **271**, 1757–1763. (doi:10.1098/rspb.2004.2757)
- Frade PR, Bongaerts P, Winkelhagen AJ, Tonk L, Bak RPM. 2008 In situ photobiology of corals over large depth ranges: a multivariate analysis on the roles of environment, host, and algal symbiont. *Limnol. Oceanogr.* **53**, 2711. (doi:10.4319/lo.2008.53.6.2711)
- Bongaerts P, Ridgway T, Sampayo EM, Hoegh-Guldberg O. 2010 Assessing the ‘deep reef refugia’ hypothesis: focus on Caribbean reefs. *Coral Reefs* **29**, 309–327. (doi:10.1007/s00338-009-0581-x)
- LaJeunesse T. 2002 Diversity and community structure of symbiotic dinoflagellates from Caribbean coral reefs. *Mar. Biol.* **141**, 387–400. (doi:10.1007/s00227-002-0829-2)
- LaJeunesse TC, Loh WK, van Woessik R, Hoegh-Guldberg O, Schmidt GW, Fitt WK. 2003 Low symbiont diversity in southern Great Barrier Reef corals, relative to those of the Caribbean. *Limnol. Oceanogr.* **48**, 2046–2054. (doi:10.4319/lo.2003.48.5.2046)
- LaJeunesse TC, Pettay DT, Sampayo EM, Phongsuwan N, Brown B, Obura DO, Hoegh-Guldberg O, Fitt WK. 2010 Long-standing environmental conditions, geographic isolation and host-symbiont specificity influence the relative ecological dominance and genetic diversification of coral endosymbionts in the genus *Symbiodinium*. *J. Biogeogr.* **37**, 785–800. (doi:10.1111/j.1365-2699.2010.02273.x)
- Finney JC, Pettay DT, Sampayo EM, Warner ME, Oxenford HA, LaJeunesse TC. 2010 The relative significance of host-habitat, depth, and geography on the ecology, endemism, and speciation of coral endosymbionts in the genus *Symbiodinium*. *Microb. Ecol.* **60**, 250–263. (doi:10.1007/s00248-010-9681-y)
- Cooper TF, Ulstrup KE, Dandan SS, Heyward AJ, Kühl M, Muirhead A, O’Leary RA, Ziersen BE, Van Oppen MJ. 2011 Niche specialization of reef-building corals in the mesophotic zone: metabolic trade-offs between divergent *Symbiodinium* types. *Proc. R. Soc. B* **278**, 1840–1850. (doi:10.1098/rspb.2010.2321)
- van Oppen MJ, Bongaerts P, Underwood JN, Peplow LM, Cooper TF. 2011 The role of deep reefs in shallow reef recovery: an assessment of vertical connectivity in a brooding coral from west and east Australia. *Mol. Ecol.* **20**, 1647–1660. (doi:10.1111/j.1365-294X.2011.05050.x)
- Bongaerts P *et al.* 2013 Sharing the slope: depth partitioning of agaricid corals and associated *Symbiodinium* across shallow and mesophotic habitats (2–60 m) on a Caribbean reef. *BMC Evol. Biol.* **13**, 205. (doi:10.1186/1471-2148-13-205)
- Richmond RH, Hunter CL. 1990 Reproduction and recruitment of corals: comparisons among the Caribbean, the tropical Pacific, and the Red Sea. *Mar. Ecol. Prog. Ser.* **60**, 185–203. (doi:10.3354/meps060185)
- Baird AH, Guest JR, Willis BL. 2009 Systematic and biogeographical patterns in the reproductive biology of scleractinian corals. *Annu. Rev. Ecol. Syst.* **40**, 551–571. (doi:10.1146/annurev.ecolsys.110308.120220)
- Bongaerts P, Sampayo EM, Bridge TC, Ridgway T, Vermeulen F, Englebert N, Webster JM, Hoegh-Guldberg O. 2011 *Symbiodinium* diversity in

- mesophotic coral communities on the Great Barrier Reef: a first assessment. *Mar. Ecol. Prog. Ser.* **439**, 117–126. (doi:10.3354/meps09315)
28. Chan YL, Pochon X, Fisher MA, Wagner D, Concepcion GT, Kahng SE, Toonen RJ, Gates RD. 2009 Generalist dinoflagellate endosymbionts and host genotype diversity detected from mesophotic (67–100m depths) coral *Leptoseris*. *BMC Ecol.* **9**, 21. (doi:10.1186/1472-6785-9-21)
 29. Kahng E, Copus M, Wagner D. 2014 Recent advances in the ecology of mesophotic coral ecosystems (MCEs). *Curr. Opin. Environ. Sustain.* **7**, 72–81. (doi:10.1016/j.cosust.2013.11.019)
 30. Slattery M, Lesser MP, Brazeau D, Stokes MD, Leichter JJ. 2011 Connectivity and stability of mesophotic coral reefs. *J. Exp. Mar. Biol. Ecol.* **408**, 32–41. (doi:10.1016/j.jembe.2011.07.024)
 31. Bongaerts P, Riginos C, Ridgway T, Sampayo EM, van Oppen MJ, Englebert N, Vermeulen F, Hoegh-Guldberg O. 2010 Genetic divergence across habitats in the widespread coral *Seriatopora hystrix* and its associated *Symbiodinium*. *PLoS ONE* **5**, e10871. (doi:10.1371/journal.pone.0010871)
 32. Frade PR, Reyes-Nivia MC, Faria J, Kaandorp JA, Luttikhuisen PC, Bak RP. 2010 Semi-permeable species boundaries in the coral genus *Madracis*: introgression in a brooding coral system. *Mol. Phylogenet. Evol.* **57**, 1072–1090. (doi:10.1016/j.ympev.2010.09.010)
 33. Brazeau DA, Lesser MP, Slattery M. 2013 Genetic structure in the coral, *Montastraea cavernosa*: assessing genetic differentiation among and within mesophotic reefs. *PLoS ONE* **8**, e65845. (doi:10.1371/journal.pone.0065845)
 34. Serrano X, Baums IB, O'Reilly K, Smith TB, Jones RJ, Shearer TL, Nunes FLD, Baker AC. 2014 Geographic differences in vertical connectivity in the Caribbean coral *Montastraea cavernosa* despite high levels of horizontal connectivity at shallow depths. *Mol. Ecol.* **23**, 4226–4240. (doi:10.1111/mec.12861)
 35. Shearer TL, Gutiérrez-Rodríguez C, Coffroth MA. 2005 Generating molecular markers from zooxanthellate cnidarians. *Coral Reefs* **24**, 57–66. (doi:10.1007/s00338-004-0442-6)
 36. Wilson K *et al.* 2002 Genetic mapping of the black tiger shrimp *Penaeus monodon* with amplified fragment length polymorphism. *Aquaculture* **204**, 297–309. (doi:10.1016/S0044-8486(01)00842-0)
 37. Clement M, Posada D, Crandall KA. 2000 TCS: a computer program to estimate gene genealogies. *Mol. Ecol.* **9**, 1657–1659. (doi:10.1046/j.1365-294x.2000.01020.x)
 38. Franklin EC, Stat M, Pochon X, Putnam HM, Gates RD. 2012 GeoSymbio: a hybrid, cloud-based web application of global geospatial bioinformatics and eoinformatics for *Symbiodinium*–host symbioses. *Mol. Ecol. Res.* **12**, 369–373. (doi:10.1111/j.1755-0998.2011.03081.x)
 39. Santos RS, LaJeunesse TC. 2006 Searchable database of symbiodinium diversity–geographic and ecological diversity (SD2-GED). Auburn, AL: Auburn University. See www.auburn.edu/~santosr/sd2_ged.htm.
 40. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013 MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* **30**, 2725–2729. (doi:10.1093/molbev/mst197)
 41. Baker AC, Rowan R, Knowlton N. 1997 Symbiosis ecology of two Caribbean acroporid corals. In *Proc. 8th Int. Coral Reef Symp.* **2**, 1295–1300.
 42. Toller WW, Rowan R, Knowlton N. 2001 Zooxanthellae of the *Montastraea annularis* species complex: patterns of distribution of four taxa of *Symbiodinium* on different reefs and across depths. *Biol. Bull.* **201**, 348–359. (doi:10.2307/1543613)
 43. Tonk L, Sampayo EM, LaJeunesse TC, Schrammeyer V, Hoegh-Guldberg O. 2014 *Symbiodinium* (Dinophyceae) diversity in reef-invertebrates along an offshore to inshore reef gradient near lizard island, great barrier reef. *J. Phycol.* **50**, 552–563. (doi:10.1111/jpy.12185)
 44. Rowan R. 2004 Coral bleaching: thermal adaptation in reef coral symbionts. *Nature* **430**, 742. (doi:10.1038/430742a)
 45. Tchernov D, Gorbunov MY, de Vargas C, Narayan Yadav S, Milligan AJ, Häggblom M, Falkowski PG. 2004 Membrane lipids of symbiotic algae are diagnostic of sensitivity to thermal bleaching in corals. *Proc. Natl Acad. Sci. USA* **101**, 13 531–13 535. (doi:10.1073/pnas.0402907101)
 46. Thornhill DJ, LaJeunesse TC, Kemp DW, Fitt WK, Schmidt GW. 2006 Multi-year, seasonal genotypic surveys of coral-algal symbioses reveal prevalent stability or post-bleaching reversion. *Mar. Biol.* **148**, 711–722. (doi:10.1007/s00227-005-0114-2)
 47. Willis BL, van Oppen MJ, Miller DJ, Vollmer SV, Ayre DJ. 2006 The role of hybridization in the evolution of reef corals. *Annu. Rev. Ecol. Syst.* **37**, 489–517. (doi:10.1146/annurev.ecolsys.37.091305.110136)
 48. Loh WKW, Loi T, Carter D, Hoegh-Guldberg O. 2001 Genetic variability of the symbiotic dinoflagellates from the wide ranging coral species *Seriatopora hystrix* and *Acropora longicyathus* in the Indo-West Pacific. *Mar. Ecol. Prog. Ser.* **222**, 97–107. (doi:10.3354/meps222097)
 49. LaJeunesse TC, Thornhill DJ, Cox EF, Stanton FG, Fitt WK, Schmidt GW. 2004 High diversity and host specificity observed among symbiotic dinoflagellates in reef coral communities from Hawaii. *Coral Reefs* **23**, 596–603.
 50. Byler KA, Carmi-Veal M, Fine M, Goulet TL. 2013 Multiple symbiont acquisition strategies as an adaptive mechanism in the coral *Stylophora pistillata*. *PLoS ONE* **8**, e59596. (doi:10.1371/journal.pone.0059596)
 51. van Oppen JH. 2004 Mode of zooxanthella transmission does not affect zooxanthella diversity in acroporid corals. *Mar. Biol.* **144**, 1–7. (doi:10.1007/s00227-003-1187-4)
 52. Stat M, Loh WKW, Hoegh-Guldberg O, Carter DA. 2008 Symbiont acquisition strategy drives host–symbiont associations in the southern Great Barrier Reef. *Coral Reefs* **27**, 763–772. (doi:10.1007/s00338-008-0412-5)
 53. Schmidt-Roach S, Miller KJ, Lundgren P, Andreakis N. 2014 With eyes wide open: a revision of species within and closely related to the *Pocillopora damicornis* species complex (Scleractinia; Pocilloporidae) using morphology and genetics. *Zool. J. Linn. Soc.* **170**, 1–33. (doi:10.1111/zooj.12092)
 54. Fabina NS, Putnam HM, Franklin EC, Stat M, Gates RD. 2012 Transmission mode predicts specificity and interaction patterns in coral–*Symbiodinium* networks. *PLoS ONE* **7**, e44970. (doi:10.1371/journal.pone.0044970)
 55. Chornesky EA, Peters EC. 1987 Sexual reproduction and colony growth in the scleractinian coral *Porites astreoides*. *Biol. Bull.* **172**, 161–177. (doi:10.2307/1541790)
 56. LaJeunesse TC. 2005 'Species' radiations of symbiotic dinoflagellates in the Atlantic and Indo-Pacific since the Miocene–Pliocene transition. *Mol. Biol. Evol.* **22**, 570–581. (doi:10.1093/molbev/msi042)
 57. Howells EJ, Beltran VH, Larsen NW, Bay LK, Willis BL, Van Oppen MJH. 2012 Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nat. Clim. Change* **2**, 116–120. (doi:10.1038/nclimate1330)
 58. Barshis DJ, Stillman JH, Gates RD, Toonen RJ, Smith LW, Birkeland C. 2010 Protein expression and genetic structure of the coral *Porites lobata* in an environmentally extreme Samoan back reef: does host genotype limit phenotypic plasticity? *Mol. Ecol.* **19**, 1705–1720. (doi:10.1111/j.1365-294X.2010.04574.x)
 59. Serrano X. 2013 Horizontal versus vertical connectivity in Caribbean reef corals: identifying potential sources of recruitment following disturbance. PhD thesis, University of Miami, Coral Gables, FL, USA.