

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/278035769>

First record of *Litarachna caribica* (Acari, Pontarachnidae) from the Pacific coast of Panama

Article in *Marine Biodiversity Records* · June 2015

DOI: 10.1017/S1755267215000688

CITATIONS

0

READS

52

3 authors:



Vladimir Pešić

University of Montenegro

344 PUBLICATIONS 1,781 CITATIONS

SEE PROFILE



Tapas Chatterjee

Principal, Crescent International School, Dha...

146 PUBLICATIONS 551 CITATIONS

SEE PROFILE



Nikolaos V. Schizas

University of Puerto Rico at Mayagüez

85 PUBLICATIONS 775 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Faunistic Diversity of Sarnena Gora Mountains [View project](#)



fauna of terrestrial Parasitengona of Iran and Montenegro [View project](#)

All content following this page was uploaded by [Nikolaos V. Schizas](#) on 12 June 2015.

The user has requested enhancement of the downloaded file.

First record of *Litarachna caribica* (Acari, Pontarachnidae) from the Pacific coast of Panama

VLADIMIR PEŠIĆ¹, TAPAS CHATTERJEE² AND NIKOLAOS V. SCHIZAS³

¹Department of Biology, University of Montenegro, Cetinjski put b.b., 81000 Podgorica, Montenegro, ²Department of Biology, Indian School of Learning, I.S.M. Annexe, P.O. – I.S.M., Dhanbad-826004, Jharkhand, India, ³Department of Marine Sciences, University of Puerto Rico, Mayagüez, Call Box 9000, Mayagüez, PR 00681, USA

We documented the existence of a population of the southern Caribbean pontarachnid mite Litarachna caribica for the first time on the Pacific coast of Panama. Based on morphological observations, this is the first record of a pontarachnid mite with a trans-isthmian distribution, which can be explained by either modern biological dispersal or historical vicariance hypotheses. Litarachna caribica had either passed through the Panama Canal, successfully colonizing the opposite coast, or previously continuously distributed populations had become disjunct after the rise of the Central American land.

Keywords: water mites, new record, trans-isthmian distribution, range expansion

Submitted 8 April 2015; accepted 14 May 2015

INTRODUCTION

The water mite family Pontarachnidae Koenike, 1910, is the only family of the Hydrachnidia occurring in the marine environment. The family represents a well-defined monophyletic clade, but almost nothing is known about the distribution and the life cycle of Pontarachnidae (Pešić *et al.*, 2012a). Knowledge on the pontarachnid fauna from the Caribbean region has increased substantially in recent years (Pešić *et al.*, 2008, 2102b, 2014). So far, four species are known from the Caribbean Sea: *Litarachna degiustii* Cook, 1958, *Litarachna caribica* Pešić, Chatterjee & Schizas, 2008, *Litarachna lopezae* Pešić, Chatterjee, Alfaro & Schizas, 2014 and *Pontarachna nemethi* Pešić, Chatterjee & Schizas, 2012. However, an understanding of the species composition, distribution, and biogeographic and ecologic relationships of the Caribbean pontarachnid fauna is not yet possible.

This paper is focused on *Litarachna caribica* recorded for the first time from the Pacific coast of Panama. *Litarachna caribica* was described by Pešić *et al.* (2008) from Curaçao, Netherlands Antilles (Figure 1). No further record of this species was published, which limits our understanding of its distribution in the southern Caribbean. The pontarachnid fauna of the Pacific coast of the American continents is not well known. At present, only one species, *Pontarachna cruciata* Hall, 1912 from California (Hall, 1912), has been described.

MATERIAL AND METHODS

Substrata samples (algae, rubble) were collected while snorkelling off the coast of Taboguilla Island, Panama, 8°48'5.90"N 79°30'52.19"W, 4 m depth, 3 February 2009. All material was washed through a 500 and 63 µm sieve and the separated fauna were preserved in ethanol. Small metazoans were sorted under a dissecting microscope and pontarachnid mites were isolated in vials filled with ethanol. In total, two males and five deutonymphs of *Litarachna caribica* Pešić *et al.*, 2008 were collected. Collection permits were secured through the Smithsonian Tropical Research Institute (STRI), Panama City. Slide-mounting was done in Hoyer's fluid and specimens were deposited in the collection belonging to the senior author.

All measurements provided in the results section are given in µm. The following abbreviations are used: Cx-I = first coxae, dL = dorsal length, H = height, L = length, I-Leg-6 = Leg 1, sixth segment (tarsus), P-1 = palp, first segment, W = width.

RESULTS AND DISCUSSION

The specimens examined in this study agree well with the description of *Litarachna caribica* from the southern Caribbean (Pešić *et al.*, 2008). This species is characterized by the following features: (1) Cx-I medially a partially fused (in adults, fused lightly at the posterior margin of gphantosomal bay, in deutonymphs, fusion more enlarged but not exceeding half of the median length of Cx-I); (2) glandularium-like structure posterior Cx-IV not fused with adjoining coxoglandularia 4 (Figure 2, arrow 1); (3) 45–54 pairs of perigenital setae around male genital field (Figure 2, arrow 2). Although *Litarachna caribica* is similar to *Litarachna degiustii* Cook,

Corresponding author:
V. Pešić
Email: vladopesic@gmail.com




Fig. 1. Recorded distribution (marked with asterisk) of *Litarachna caribica* Pešić et al., 2008.

1958, and can coexist in the southern Caribbean, the latter can be distinguished by the glandularium-like structure posterior to Cx-IV fused with adjoining coxoglandularia 4, Cx-I medially broadly fused and in males only three pairs of perigenital setae lying around the genital field (Pešić et al., 2008).

Here, we give some measurements of the male and deutonymph from the island of Taboguilla. Male: Idiosoma L 290–341; coxal field: L 129, Cx-III W 226; genital field L/W 28/25,

48–52 perigenital setae around genital field; palp: total L 198; dL/H (in parentheses dL/H ratio): P-1, 15/19 (0.83); P-2, 60/31 (1.94); P-3, 25/28 (0.9); P-4, 72/19 (3.8); P-5, 26/12 (2.3); dorsal length of I-Leg-3–6: 38, 42, 60, 80.

Deutonymph: Idiosoma L/W 228–237/188; coxal field: L 83, Cx-III W 120; palp: total L 140; dL/H (in parentheses dL/H ratio): P-1, 10/12 (0.8); P-2, 41/20 (2.0); P-3, 19/19 (1.0); P-4, 49/14 (3.6); P-5, 21/8 (2.7).

This is the first record of a pontarachnid mite with a trans-isthmian distribution. If the species passed through the Panama Canal and successfully colonized the opposite coast, this will be the first time such an isthmian transfer has been reported. The Panama Canal represents a potential pathway for the interoceanic dispersal of marine organisms (Ruiz et al., 2009). However, many studies (Hildebrand, 1939; McCosker & Dawson, 1975; Davidson et al., 2008; Ros et al., 2014) have documented that some organisms can survive the passage through the Canal, in most cases associated with fouling on ships or in ballast water.

The trans-isthmian distribution of *Litarachna caribica* is difficult to explain without testing its potential for surviving freshwater immersion during passage through the Canal. The two freshwater lakes (lakes Gatun and Miraflores) that the boats and marine organisms have to cross, to travel from the Caribbean Sea to the Pacific Ocean, effectively serve as a barrier for most marine species. However, some pontarachnid species have been described from freshwaters (Cook, 1996), although, these species were also found to occur in estuaries or locations near the sea (Smit, 2009). The influence of the salinity in the particular case of *Litarachna denhami* has been documented by Witte & Olomski (1999), which showed that the species is a very effective hyporegulator. Some pontarachnid species are euryhaline and preadapted




Fig. 2. Photograph of *Litarachna caribica* Pešić et al., 2008, male, Taboguilla Island, Panama: coxal and genital field (arrow 1 showing a glandularium-like structure not fused with a platelet with coxoglandularia 4 and associated seta, arrow 2 showing perigenital setae around genital field).

to successfully colonize brackish water areas. However, despite the possibility that *Litarachna caribica* has survived the crossing of the Panama Canal, it is not possible to assert whether this species has recently been introduced into the Pacific or if the trans-isthmian distribution is a result of vicariance through the rise of the Panamanian Isthmus. In the late Pliocene, 3.1 million years ago, the rise of the Central American land strip created disjunct populations of previously continuously distributed species. Many studies (Knowlton *et al.*, 1993; Knowlton & Weigt, 1998; Lessios, 2008) examined the genetic divergence of 'sibling species' or 'geminate species' with trans-isthmian distribution, showing the effects of the best-known biogeographic barrier on shallow water marine fauna. Similarly, *Litarachna caribica* should be assessed with molecular markers to test whether the morphologically identical trans-isthmian mite populations are genetically similar, indicating a recent dispersal process, or if they are genetically divergent, indicating an ancient separation.

ACKNOWLEDGEMENTS

We thank the undergraduate students Mairim Ramirez Cruz, Yisenia Nazario Pabon and Rocio Vergara Torres for sorting the sediments and extracting the fauna.

REFERENCES

- Cook D.R. (1996) A freshwater species of *Pontarachna* (Acari, Pontarachnidae) from South Africa, with a discussion of genital acetabula in the family. *Anales Instituto de Biología, Universidad Nacional Autónoma de México, Seria Zoología* 67, 259–264.
- Davidson I.C., McCann L.D., Fofonoff P.W., Sytsma M.D. and Ruiz G.M. (2008) The potential for hull-mediated species transfers by obsolete ships on their final voyages. *Diversity and Distributions* 14, 518–529.
- Hall H.V.M. (1912) Some marine and terrestrial acarina of Laguna Beach. *First annual Report of Laguna Marine Laboratory, Pomona College, Claremont, California*, 1, pp. 117–186.
- Hildebrand S.F. (1939) The Panama Canal as a passageway for fishes, with lists and remarks on the fishes and invertebrates observed. *Zoologica* 24, 15–45.
- Knowlton N. and Weigt L.A. (1998) New dates and new rates for divergence across the Isthmus of Panama. *Proceedings of the Royal Society B: Biological Sciences* 265, 2257–2263.
- Knowlton N., Weigt L.A., Solórzano L.A., Mills D.K. and Bermingham E. (1993) Divergence in proteins, mitochondrial DNA, and reproductive compatibility across the Isthmus of Panama. *Science* 260, 1629–1632.
- Lessios H.A. (2008) The great American schism: divergence of marine organisms after the rise of the Central American Isthmus. *Annual Review of Ecology, Evolution, and Systematics* 39, 63–91.
- McCosker J.E. and Dawson C.E. (1975) Biotic passage through the Panama Canal, with particular reference to fishes. *Marine Biology* 30, 343–351.
- Pešić V., Chatterjee T., Alfaro M. and Schizas N.V. (2014) A new species of *Litarachna* (Acari, Hydrachnida, Pontarachnidae) from a Caribbean mesophotic coral ecosystem. *ZooKeys* 425, 89–97.
- Pešić V., Chatterjee T., Ingole B., Velip D. and Pavičević A. (2012a) A new species of *Litarachna* Walter, 1925 (Acari: Hydrachnida) from the West Indian Coast, with a discussion on the diversity of the family Pontarachnidae Koenike, 1910. *Cahiers de Biologie Marine* 53, 547–553.
- Pešić V., Chatterjee T. and Schizas N.V. (2008) Marine water mites (Acari: Hydrachnida: Pontarachnidae) from the Caribbean Sea, with description of one new species. *Cahiers de Biologie Marine* 49, 253–259.
- Pešić V., Chatterjee T. and Schizas N.V. (2012b) A new species of *Pontarachna* (Acari, Hydrachnida, Pontarachnidae) from a mesophotic coral ecosystem off Vieques Island, Puerto Rico, Caribbean Sea. *Zootaxa* 3440, 63–67.
- Ros M., Ashton G., Lacerda M., Carlton J.T., Vázquez-Luis M., Guerra-García J.M. and Ruiz G.M. (2014) The Panama Canal and the transoceanic dispersal of marine invertebrates: evaluation of the introduced amphipod *Paracaprella pusilla* Mayer, 1890 in the Pacific Ocean. *Marine Environmental Research* 99, 204–211.
- Ruiz G.M., Torchin M.E. and Grant K. (2009) Using the Panama Canal to test predictions about tropical marine invasions. *Smithsonian Contributions to the Marine Sciences* 38, 291–300.
- Smit H. (2009) Water mites of the family Pontarachnidae from Singapore, with a description of one new species (Acari: Hydrachnida). *Raffles Bulletin of Zoology, Supplement* 22, 203–205.
- and
- Witte H. and Olmski R. (1999) The evolutionary transformation of functional systems in the Parasitengona. In Needham G.R., Mitchell R., Horn D.J. and Welbourn W.C. (eds) *Acarology Proceedings IX*, Vol. 2. Columbus, OH: The Ohio Biological Survey, pp. 125–137.

Correspondence should be addressed to:

V. Pešić
Department of Biology, University of Montenegro
Cetinjski put b.b., 81000 Podgorica, Montenegro
Email: vladopesic@gmail.com